arXiv:1301.3787v2 [hep-ph] 29 Jan 2013

Modulated decay in the multi-component Universe

Seishi Enomoto
Department of Physics, Nagoya University, Nagoya 464-8602, Japan

Kazunori Kohri
Cosmophysics group, Theory Center, IPNS, KEK,
and The Graduate University for Advanced Study (Sokendai), Tsukuba 305-0801, Japan

Tomohiro Matsuda
Laboratory of Physics, Saitama Institute of Technology, Fukaya, Saitama 369-0293, Japan

The early Universe may have oscillations, kinations (nonoscillatory evolution), topological defects, relativistic and non-relativistic particles at the same time. These components, which may have distinguishable density scalings, may decay modulated. In this paper we study generation of the curvature perturbations caused by the modulated decay in the multi-component Universe. For our analytic calculation we consider the instant-decay approximation. The non-Gaussianity parameter is separated into the model-dependent and independent contributions, where the sign of the model-independent contribution is determined by the density scalings. We also find that the component, whose scaling is very close to the radiation, can generate significant non-Gaussianity from the model-independent part, and its sign could be either positive or negative. Importantly, this result shows that significant non-Gaussianity can be generated even if the decaying component is already dominating the universe. We also consider partial decay, in which a fraction of the component decays modulated.

PACS numbers: 98.80Cq

I. INTRODUCTION

Our focus in this paper is the primordial curvature perturbation $\zeta(k)$ that exists on cosmological scales just before they start to enter the horizon. The cosmological observations are strongly constraining those cosmological perturbations and providing a unique window on the very early Universe [1].

Generation of the cosmological perturbations begins presumably during inflation, when the vacuum fluctuations of bosonic fields are converted to a classical perturbation. Within this general framework, one can find many proposals [1–10, 12, 13].

First recall the δN formalism used to calculate ζ. To define the curvature perturbation ζ, the energy density ρ is smoothed on a super-horizon scale shorter than any scale of interest. Then the local energy continuity equation is given by

$$\frac{\partial\rho(x,t)}{\partial t} = -\frac{3}{a(x,t)}\frac{\partial a(x,t)}{\partial t}\left(\rho(x,t)+p(x,t)\right), \qquad (1)$$

where t is time along a comoving thread of spacetime and $a(t)$ is the local scale factor. During nearly exponential inflation, the vacuum fluctuation of each light scalar field ϕ_i is converted at horizon exit to a nearly Gaussian classical perturbation with spectrum $(H/2\pi)^2$, where $H \equiv \dot{a}(t)/a(t)$ in the unperturbed Universe. Writing the curvature perturbation

$$\zeta = \delta[\ln(a(x,t)/a(t_1)] \equiv \delta N, \qquad (2)$$

and taking t_* to be an epoch during inflation after relevant scales leave the horizon, we assume $N(\phi_1(x,t_*), \phi_2(x,t_*), \cdots, t, t_*)$ so that

$$\zeta(x,t) = N_i\delta\phi_i(x,t_*) + \frac{1}{2}N_{ij}\delta\phi_j(x,t_*) + \cdots, \qquad (3)$$

where a subscript i denotes $\partial/\partial\phi_i$ evaluated on the unperturbed trajectory. The δN-formalism is also valid for the evolution after inflation.

We consider modulation of the decay rate $\Gamma(\varphi)$, which causes inhomogeneous decay of the component (ρ_σ) in the multi-component Universe. We are assuming instant decay for the calculation [9]. See Fig.1 for the basic setups of the modulated reheating scenario.

The source of the modulation is the moduli perturbation caused by an additional light field φ, whose potential is assumed to be negligible at the decay. The "seed" perturbation $\delta\varphi$ is generated during the primordial inflation. At the horizon exit, we consider Gaussian perturbation $\delta\varphi_* \equiv \varphi_* - \bar{\varphi}_*$. At the decay, we introduce the function $\varphi_d = g(\varphi_*)$ and the expansion about the Gaussian perturbation $\delta\varphi \equiv g'\delta\varphi_*$ (or equivalently $\delta\varphi_* = \delta\varphi/g'$). The function g explains evolution after the horizon exit [16].

In our scenario of the multi-component Universe, we assume that the component ρ_σ decays after reheating.[1] For the first example, we consider simple two-component

---

[1] In the multi-component Universe, the calculation may generically depend on isocurvature perturbations between components. Although we are calculating the perturbations when the isocurvature perturbations are negligible, our formalism is carefully prepared so that the mixed perturbations can be calculated within the formalism.



FIG. 1: Modulation at the transition causes density perturbations when the decaying component changes its density scaling. The straight line shows the instant-decay approximation.

Universe, which has $\rho_\sigma \propto a^{-3}$ (matter) and $\rho_r \propto a^{-4}$ (radiation) before the decay.

Later in this paper we are going to extend our analytic calculation to the components that may "not" scale like matter [13–15]; typical examples are the cosmological defects or the oscillatory / nonoscillatory evolutions [14] that may be caused by the non-quadratic potentials.

II. MODULATED DECAY IN THE SIMPLE MULTI-COMPONENT UNIVERSE

First consider the simple (matter + radiation) multi-component Universe. The set-ups are similar to the curvaton. In the curvaton mechanism [5], the significant contribution comes from the evolution **before** the decay; while the modulated decay [9] describes the generation of the curvature perturbations **at the end** [10, 11].

For our analytic calculation, we consider instant-decay approximation [9]. The actual transition could be more complicated depending on the details of the model parameters. For the modulated reheating scenario, the idea of the continual decay has been considered by many authors [17, 18]. In fact, the Universe in which matter is decaying continually into radiation can realize the multi-component Universe during the decay. In that way, just by shifting the initial conditions, previous numerical studies could have been including "the modulated decay in the multi-component Universe". In that sense numerical calculation is versatile. On the other hand, the analytic estimation, which is calculated in specific limits, will help understand the physics behind it. In our analytic calculation we found that model-independent non-Gaussianity is significant both in the single and in the multi-component Universe. The model-dependent non-Gaussianity is calculated from the perturbations of $\Gamma(\varphi_*)$. Curvature perturbation in models where a light scalar field is the curvaton and the moduli that induces modulated reheating the same time has been considered in Ref.[19]. We hope the reader can easily compare our calculation with those previous studies.

In our model, the uniform density hypersurfaces defined for the instant decay is given by



FIG. 2: In the left picture we show the densities and their scalings in the modulated ($\Gamma \neq \Gamma_0$) patch. The right picture shows the unmodulated (reference) Universe. These pictures are sharing Γ_0 ($\rho_{\sigma,0}$ and $\rho_{r,0}$) and Γ as the common quantities. Due to the different ρ_σ-scalings after Γ_0, one will find $N_m \neq N_{\not{m}}$ and $\rho_{i,\Gamma,m} \neq \rho_{i,\Gamma,\not{m}}$.

$$\rho_{\sigma,\Gamma} + \rho_{r,\Gamma} \;\equiv\; 3M_p^2\Gamma^2, \tag{4}$$

where the instant decay occurs at $H=\Gamma$. More specifically, in the modulated Universe (see the left picture in Fig.2) we have

$$\rho_{\sigma,\Gamma,m} + \rho_{r,\Gamma,m} \;\equiv\; 3M_p^2\Gamma^2, \tag{5}$$

and in the unmodulated (reference) Universe (see the right picture in Fig.2) we have

$$\rho_{\sigma,\Gamma,\not{m}} + \rho_{r,\Gamma,\not{m}} \;\equiv\; 3M_p^2\Gamma^2. \tag{6}$$

Introducing the mean value Γ_0, we define the uniform density hypersurface

$$\rho_{\sigma,0} + \rho_{r,0} \;\equiv\; 3M_p^2\Gamma_0^2, \tag{7}$$

which is common in the modulated and in the unmodulated Universe. Without loss of generality, one may choose $\Gamma < \Gamma_0$ for the calculation.

Using the density scalings, we find in the modulated Universe

$$\begin{aligned}
\rho_{\sigma,\Gamma,m} &= \rho_{\sigma,0}\left(\frac{a_{\Gamma,m}}{a_{\Gamma_0}}\right)^{-3} \\
\rho_{r,\Gamma,m} &= \rho_{r,0}\left(\frac{a_{\Gamma,m}}{a_{\Gamma_0}}\right)^{-4},
\end{aligned} \tag{8}$$

which lead to

$$\frac{\rho_{\sigma,0}\left(\frac{a_{\Gamma,m}}{a_{\Gamma_0}}\right)^{-3} + \rho_{r,0}\left(\frac{a_{\Gamma,m}}{a_{\Gamma_0}}\right)^{-4}}{\rho_{\sigma,0}+\rho_{r,0}} = \frac{\Gamma^2}{\Gamma_0^2}. \tag{9}$$

Defining local "N_m" by

$$N_m \equiv \int_{t_{\Gamma_0}}^{t_{\Gamma,m}} H(t)dt, \tag{10}$$

where $H(t) \equiv \dot{a}/a$, one can rewrite the above equation as

$$f_\sigma e^{-3N_m} + (1-f_\sigma)e^{-4N_m} \;=\; \frac{\Gamma^2}{\Gamma_0^2}, \tag{11}$$

where the coefficient is defined by

$$f_\sigma \;\equiv\; \frac{\rho_{\sigma,0}}{\rho_{\sigma,0}+\rho_{r,0}}. \tag{12}$$

In order to compare N_m with the unmodulated Universe, we find the local equation in the unmodulated Universe,

$$f_\sigma e^{-4N_{\not{m}}} + (1-f_\sigma)e^{-4N_{\not{m}}} \;=\; \frac{\Gamma^2}{\Gamma_0^2}. \tag{13}$$

Let us include isocurvature perturbations at $H=\Gamma_0$. We consider the non-linear formalism [20, 21] applied just before $H=\Gamma_0$;

$$\begin{aligned}
\zeta_\sigma &= \delta N_{\rm ini} + \frac{1}{3}\int_{\bar\rho_{\sigma,0}}^{\rho_{\sigma,0}} \frac{d\tilde\rho_\sigma}{\tilde\rho_\sigma} && (14)\\
\zeta_r &= \delta N_{\rm ini} + \frac{1}{4}\int_{\bar\rho_{r,0}}^{\rho_{r,0}} \frac{d\tilde\rho_r}{\tilde\rho_r}, && (15)
\end{aligned}$$

where $\rho_{\sigma,0}$ and $\rho_{r,0}$ are defined on the uniform density hypersurfaces, and $\bar\rho_i$ denotes their mean value. Here $\delta N_{\rm ini}$ denotes the curvature perturbation at that moment, which could be significant if the curvaton evolution before the decay is significant.

Finally, the non-linear formalism gives

$$\begin{aligned}
\rho_{\sigma,0} &= \bar\rho_{\sigma,0}e^{3(\zeta_\sigma-\delta N_{\rm ini})} && (16)\\
\rho_{r,0} &= \bar\rho_{\sigma,0}e^{4(\zeta_r-\delta N_{\rm ini})}, && (17)
\end{aligned}$$

which leads to

$$\begin{aligned}
f_\sigma &= \frac{\bar\rho_{\sigma,0}e^{3(\zeta_\sigma-\delta N_{\rm ini})}}{\bar\rho_{\sigma,0}e^{3(\zeta_\sigma-\delta N_{\rm ini})}+\bar\rho_{r,0}e^{4(\zeta_r-\delta N_{\rm ini})}}\\
&= \frac{\bar\rho_{\sigma,0}e^{3(\zeta_\sigma-\delta N_{\rm ini})}}{3M_p^2\Gamma_0^2}. && (18)
\end{aligned}$$

Using the component perturbations ζ_σ and ζ_r, we find

$$\begin{aligned}
\bar f_\sigma e^{3(\zeta_\sigma-\delta N_{\rm ini}-N_m)} + (1-\bar f_\sigma)e^{4(\zeta_r-\delta N_{\rm ini}-N_m)} &= \frac{\Gamma^2}{\Gamma_0^2}\\
\bar f_\sigma e^{3(\zeta_\sigma-\delta N_{\rm ini})-4N_{\not{m}}} + (1-\bar f_\sigma)e^{4(\zeta_r-\delta N_{\rm ini}-N_{\not{m}})} &= \frac{\Gamma^2}{\Gamma_0^2},
\end{aligned} \tag{19}$$

where the coefficient is defined by

$$\bar f_\sigma \;\equiv\; \frac{\bar\rho_{\sigma,0}}{3M_p^2\Gamma_0^2}. \tag{20}$$

A. First order

From Eq.(19), we find at first order

$$\begin{aligned}
2\frac{\delta\Gamma_m^{(1)}}{\Gamma_0} &= 3\bar f_\sigma(\zeta_\sigma^{(1)}-\delta N_{\rm ini}^{(1)}-N_m^{(1)})\\
&\quad +4(1-\bar f_\sigma)(\zeta_r^{(1)}-\delta N_{\rm ini}^{(1)}-N_m^{(1)}) && (21)\\
&\quad -9\bar f_\sigma N_m^{(0)}(\zeta_\sigma^{(1)}-\delta N_{\rm ini}^{(1)})\\
&\quad -16(1-\bar f_\sigma)N_m^{(0)}(\zeta_r^{(1)}-\delta N_{\rm ini}^{(1)}) && (22)\\
2\frac{\delta\Gamma_{\not{m}}^{(1)}}{\Gamma_0} &= \bar f_\sigma(3\zeta_\sigma^{(1)}-3\delta N_{\rm ini}^{(1)}-4N_{\not{m}}^{(1)})\\
&\quad +4(1-\bar f_\sigma)(\zeta_r^{(1)}-\delta N_{\rm ini}^{(1)}-N_{\not{m}}^{(1)})\\
&\quad -12\bar f_\sigma N_{\not{m}}^{(0)}(\zeta_\sigma^{(1)}-\delta N_{\rm ini}^{(1)})\\
&\quad -16(1-\bar f_\sigma)N_{\not{m}}^{(0)}(\zeta_r^{(1)}-\delta N_{\rm ini}^{(1)}), && (23)
\end{aligned}$$

where the evolution of the perturbations between $H=\Gamma_0$ and $H=\Gamma$ vanishes at this order, because we have $N_m^{(0)}=N_{\not{m}}^{(0)}=0$.

Solving the above equations, we find

$$\begin{aligned}
N_m^{(1)} &= -p_\sigma\frac{\delta\Gamma_m^{(1)}}{\Gamma_0}+r_\sigma(\zeta_\sigma^{(1)}-\delta N_{\rm ini}^{(1)})\\
&\quad +(1-r_\sigma)(\zeta_r^{(1)}-\delta N_{\rm ini}^{(1)}) && (24)\\
N_{\not{m}}^{(1)} &= -\frac{1}{2}\frac{\delta\Gamma_{\not{m}}^{(1)}}{\Gamma_0}+\frac{3}{4}\bar f_\sigma(\zeta_\sigma^{(1)}-\delta N_{\rm ini}^{(1)})\\
&\quad +(1-\bar f_\sigma)(\zeta_r^{(1)}-\delta N_{\rm ini}^{(1)}), && (25)
\end{aligned}$$

where the coefficients are defined by

$$\begin{aligned}
p_\sigma &\equiv \frac{2(\bar\rho_{\sigma,0}+\bar\rho_{r,0})}{3\bar\rho_{\sigma,0}+4\bar\rho_{r,0}}\\
r_\sigma &\equiv \frac{3\bar\rho_{\sigma,0}}{3\bar\rho_{\sigma,0}+4\bar\rho_{r,0}}. && (26)
\end{aligned}$$

We thus find the relation

$$p_\sigma - \frac{1}{2} = \frac{1}{6}r_\sigma. \tag{27}$$

We are calculating $N_{\not{m}}$ to see the deviation from the unmodulated Universe. $N_{\not{m}}^{(1)} \neq 0$ is not the generation of the curvature perturbations in the unmodulated Universe.

Therefore, the curvature perturbation created by the modulation ($\delta N^{(1)} \equiv N_m^{(1)} - N_{\not{m}}^{(1)}$) is calculated as

$$\delta N^{(1)} \;=\; \left(-p_\sigma\frac{\delta\Gamma_m^{(1)}}{\Gamma_0}+\frac{1}{2}\frac{\delta\Gamma_{\not{m}}^{(1)}}{\Gamma_0}\right), \tag{28}$$

where the other terms cancel by definition.

To understand the above result, let us consider the perturbation of Γ. Expanding $\varphi_d \equiv g(\varphi_*)$, which defines φ at the decay, we find [22]

$$\varphi_d \;=\; \bar g + \delta\varphi. \tag{29}$$

where $\delta\varphi \equiv g'\delta\varphi_*$. In that way the first order perturbation of the decay rate is calculated as

$$\delta\Gamma^{(1)} = \left[\frac{\partial\Gamma}{\partial g}\right]_{g=\bar{g}} \delta\varphi. \tag{30}$$

In the practical calculation $\delta\Gamma_m$ and $\delta\Gamma_{\not m}$ are identical. We thus find

$$\delta N^{(1)} \;=\; \left(-p_\sigma + \frac{1}{2}\right)\frac{\delta\Gamma^{(1)}}{\Gamma_0}, \tag{31}$$

where $\delta\Gamma^{(1)} \equiv \delta\Gamma^{(1)}_m = \delta\Gamma^{(1)}_{\not m}$. In the single-component limit ($p_\sigma = 2/3$), we find

$$\begin{aligned} \delta N^{(1)} \;&=\; -\frac{1}{6}\frac{\delta\Gamma^{(1)}}{\Gamma_0} \\ &=\; -\frac{1}{6}\frac{\Gamma'}{\Gamma_0}\delta\varphi, \end{aligned} \tag{32}$$

which reproduces the calculation in Ref.[9].

It is obvious that g is trivial in the slow-roll limit; however for more practical estimation one has to calculate the model-dependent function g. We are avoiding the calculation of g, since it is highly mode-dependent.

B. Second order

Generically, one can expand

$$\varphi = \bar{\varphi} + \sum_{k=1}^{\infty}\frac{1}{k!}\delta^{(k)}\varphi, \tag{33}$$

where $\delta^{(1)}\varphi$ is a Gaussian random field. In the same way, the primordial perturbation can be expanded as

$$\zeta = \zeta^{(1)} + \sum_{k=2}^{\infty}\frac{1}{k!}\zeta^{(\mathrm{k})}, \tag{34}$$

where $\zeta^{(1)}$ is Gaussian. Non-linearity parameters are defined for the adiabatic perturbation ζ;

$$\zeta = \zeta^{(1)} + \frac{3}{5}f_{NL}(\zeta^{(1)})^2 + \frac{9}{25}g_{NL}(\zeta^{(1)})^3 + ... \tag{35}$$

Using the Gaussian quantum fluctuations at the horizon exit ($\delta\varphi_*$), we can write [22]

$$\varphi_* = \bar{\varphi}_* + \delta\varphi_*, \tag{36}$$

which is exact by definition. Again, we write

$$\varphi_d \equiv g(\varphi_*) \tag{37}$$

and expand it as [22]

$$\varphi_{\mathrm{ini}} \;=\; \bar{g} + \sum_{k=1}^{\infty}\frac{1}{k!}g^{(k)}\left(\frac{\bar{g}}{g'}\frac{\delta\varphi}{\bar{\varphi}}\right)^k, \tag{38}$$

where we wrote $g^{(k)} \equiv \partial^k g/\partial\varphi_*^k$.

1. *Decay rates*

Before discussing non-Gaussianity of the second order perturbations, let us examine the specific expansion of the model-dependent decay rates.

- Our first example is

  $$\Gamma(\varphi) = \bar{\Gamma}\left(1 + \frac{1}{2}\frac{\varphi^2}{M_*^2}\right). \tag{39}$$

  Then, one can expand

  $$\frac{\Gamma(\varphi_d)}{\bar{\Gamma}} = 1 + \frac{1}{2}\frac{\left[\bar{g} + \sum_{k=1}^{\infty}\frac{1}{k!}g^{(k)}\left(\frac{\bar{g}}{g'}\frac{\delta\varphi}{\bar{\varphi}}\right)^k\right]^2}{M_*^2}. \tag{40}$$

  We thus find for the expansion $\Gamma = \Gamma_0 + \delta\Gamma^{(1)} + \frac{1}{2}\Gamma^{(2)} + ...$ with the approximation $\bar{\Gamma} \simeq \Gamma_0$;

  $$\frac{\delta\Gamma^{(1)}}{\Gamma_0} \;\equiv\; \frac{\bar{g}}{M_*^2}\delta\varphi \tag{41}$$

  $$\begin{aligned} \frac{\delta\Gamma^{(2)}}{\Gamma_0} \;&\equiv\; \frac{1}{M_*^2}\left[1 + \frac{g''\bar{g}}{(g')^2}\right](\delta\varphi)^2 \\ &=\; \frac{M_*^2}{\bar{g}^2}\left[1 + \frac{g''\bar{g}}{(g')^2}\right]\left(\frac{\delta\Gamma^{(1)}}{\Gamma_0}\right)^2. \end{aligned} \tag{42}$$

  An interesting case would be $g \lesssim M_*$, where the initial condition is comparable but less than the cut-off scale. In that case one can find significant f_{NL} in the conceivable range. Moreover, it is possible to find negative contribution from

  $$\Gamma(\varphi) = \bar{\Gamma}\left(1 - \frac{1}{2}\frac{\varphi^2}{M_*^2}\right). \tag{43}$$

  The flip of the sign is very important.

- Second, we consider $\Gamma \propto \varphi^n$. The specific form beocmes

  $$\Gamma(\varphi) = \frac{\lambda^{(n)}}{n}\frac{\varphi^n}{M_*^{n-1}}. \tag{44}$$

  Then, Γ can be expanded as

  $$\frac{\delta\Gamma^{(1)}}{\Gamma_0} \;=\; \left[n\frac{\delta\varphi}{\bar{g}}\right] \tag{45}$$

  $$\frac{\delta\Gamma^{(2)}}{\Gamma_0} \;=\; \frac{1}{2}\left[(n-1) + \frac{\bar{g}g''}{(g')^2}\right]\left(\frac{\delta\Gamma^{(1)}}{\Gamma_0}\right)^2. \tag{46}$$

Let us summarize the results. Defining

$$\frac{\delta\Gamma^{(2)}}{\Gamma_0} \;=\; A\left(\frac{\delta\Gamma^{(1)}}{\Gamma_0}\right)^2, \tag{47}$$

we find

$$\begin{aligned} A &= \pm\frac{M_*^2}{\bar{g}^2}\left[1+\frac{g''\bar{g}}{(g')^2}\right] \\ &\quad \text{for } \Gamma(\varphi)=\Gamma_*\left(1\pm\frac{1}{2}\frac{\varphi^2}{M_*^2}\right), \end{aligned} \tag{48}$$

and

$$\begin{aligned} A &= \frac{1}{2}\left[(n-1)+\frac{\bar{g}g''}{(g')^2}\right] \\ &\quad \text{for } \Gamma(\varphi)=\frac{\lambda^{(n)}}{n}\frac{\varphi^n}{M_*^{n-1}}. \end{aligned} \tag{49}$$

The above calculations are important for the estimation of the model-dependent part of the non-Gaussianity parameter f_{NL}.

2. f_{NL}

In order to extract the contributions from the modulation, we are going to assume $\zeta_\sigma \simeq \zeta_r \simeq 0$. We also assume $\delta N_{\rm ini}\simeq 0$ for simplicity.

Then, one can easily expand Eq.(19) to find the second order perturbations. The expansions used here are

$$\begin{aligned} e^{aN} &= 1+a(N^{(1)}+\frac{1}{2}N^{(2)}+...) \\ &\quad +\frac{a^2}{2}(N^{(1)}+\frac{1}{2}N^{(2)}+...)^2+..., \end{aligned} \tag{50}$$

and

$$\left(\frac{\Gamma}{\Gamma_0}\right)^2=\left(\frac{\Gamma_0+\delta\Gamma^{(1)}+\frac{1}{2}\delta\Gamma^{(2)}+...}{\Gamma_0}\right)^2. \tag{51}$$

We find for the second order perturbations

$$\begin{aligned} N_m^{(2)} &= p_\sigma\left[\left(8-\frac{7}{2}\bar{f}_\sigma\right)\left(N_m^{(1)}\right)^2-\frac{\left(\delta\Gamma^{(1)}\right)^2+\Gamma_0\delta\Gamma^{(2)}}{\Gamma_0^2}\right] \\ N_{\not{m}}^{(2)} &= \frac{1}{2}\left[8\left(N_{\not{m}}^{(1)}\right)^2-\frac{\left(\delta\Gamma^{(1)}\right)^2+\Gamma_0\delta\Gamma^{(2)}}{\Gamma_0^2}\right]. \end{aligned} \tag{52}$$

Using the relations between the first order perturbations;

$$\begin{aligned} N_m^{(1)} &= -p_\sigma\frac{\delta\Gamma^{(1)}}{\Gamma_0} \\ &= -\frac{2p_\sigma}{1-2p_\sigma}\delta N^{(1)} \end{aligned} \tag{53}$$

$$\begin{aligned} N_{\not{m}}^{(1)} &= -\frac{1}{2}\frac{\delta\Gamma^{(1)}}{\Gamma_0} \\ &= -\frac{1}{1-2p_\sigma}\delta N^{(1)}, \end{aligned} \tag{54}$$

and the definition (47), we find

$$\begin{aligned} N_m^{(2)} &= \left[\frac{p_\sigma^3(32-14\bar{f}_\sigma)}{(1-2p_\sigma)^2}-\frac{4p_\sigma(1+A)}{(1-2p_\sigma)^2}\right]\left(\delta N^{(1)}\right)^2 \\ N_{\not{m}}^{(2)} &= \left[\frac{4}{(1-2p_\sigma)^2}-\frac{2(1+A)}{(1-2p_\sigma)^2}\right]\left(\delta N^{(1)}\right)^2. \end{aligned} \tag{55}$$

We thus find

$$\begin{aligned} f_{NL} &= \frac{5}{3}\left[\frac{p_\sigma^3(16-7\bar{f}_\sigma)-2p_\sigma-1}{(1-2p_\sigma)^2}\right] \\ &\quad +\frac{5A}{3}\frac{1}{1-2p_\sigma}, \end{aligned} \tag{56}$$

where the last term depends on A (the model-dependent contribution). In the single-component limit ($\bar{f}_\sigma\to 1$ and $p_\sigma\to 2/3$), we find a simple formula

$$f_{NL} = 5-5A. \tag{57}$$

Note that the model-independent contribution $f_{NL}=5$ is showing an interesting result. Note also that $\Gamma\propto\varphi^3$ gives $A\sim 1$ when the evolution of φ is trivial, which may lead to the cancellation ($f_{NL}\simeq 0$).

In the opposite limit, $\bar{f}_\sigma\to 0$ leads to $p_\sigma\to 1/2$. In that limit we find

$$f_{NL}\propto\frac{1}{(1-2p_\sigma)^2}\gg 1. \tag{58}$$

III. HIGHER POTENTIAL OR TOPOLOGICAL DEFECTS

For the scalar potential of the form $V(\sigma)\propto\sigma^n$, the energy density of the scalar-field oscillations decreases as $\rho_\sigma\propto a^{\frac{-6n}{n+2}}$ when the oscillations are rapid compared with the expansion rate [23]. Alternatively, one may choose topological defects for the decaying component, which may scale like $\rho_\sigma\propto a^k$. NO (Non Oscillatory) motion can lead to a different density scaling [14]. Here the scaling is approximately defined at the time of the decay; there is no need to find exact scale-dependence that is valid during the whole evolution. This point might be crucial for the practical investigation.

For our purpose, we consider the component that scales like $\rho_\sigma\propto a^{-(4+\epsilon_n)}$. Here $\epsilon_n=-1$ corresponds to the sinusoidal oscillation for the quadratic potential, whose energy density scales like $\rho_\sigma\propto a^{-3}$. Note that $\epsilon_n\geq 0$ is not excluded in our calculation; we will show that this may change the sign of f_{NL}.

In order to include the isocurvature perturbation at $H=\Gamma_0$, we consider the component perturbations defined by

$$\zeta_\sigma = \delta N_{\rm ini}+\frac{1}{4+\epsilon_n}\int_{\bar{\rho}_{\sigma,0}}^{\rho_{\sigma,0}}\frac{d\tilde{\rho}_\sigma}{\tilde{\rho}_\sigma} \tag{59}$$

$$\zeta_r = \delta N_{\rm ini}+\frac{1}{4}\int_{\bar{\rho}_{r,0}}^{\rho_{r,0}}\frac{d\tilde{\rho}_r}{\tilde{\rho}_r}. \tag{60}$$

Then we find

$$\begin{aligned}\bar{f}_\sigma e^{(4+\epsilon_n)(\zeta_\sigma-\delta N_{\rm ini}-N_m)}+(1-\bar{f}_\sigma)e^{4(\zeta_r-\delta N_{\rm ini}-N_m)} &= \frac{\Gamma^2}{\Gamma_0^2}\\ \bar{f}_\sigma e^{(4+\epsilon_n)(\zeta_\sigma-\delta N_{\rm ini})-4N_{\not{m}}}+(1-\bar{f}_\sigma)e^{4(\zeta_r-\delta N_{\rm ini}-N_{\not{m}})} &= \frac{\Gamma^2}{\Gamma_0^2}.\end{aligned} \tag{61}$$

Again, the isocurvature perturbations and $\delta N_{\rm ini}$ does not contribute to the modulation at first order. They will be neglected in the second order calculations.

A. First order

From Eq.(61), we find at first order

$$2\frac{\delta\Gamma^{(1)}_m}{\Gamma_0} = -(4+\epsilon_n)\bar{f}_\sigma N^{(1)}_m - 4(1-\bar{f}_\sigma)N^{(1)}_m \tag{62}$$

$$2\frac{\delta\Gamma^{(1)}_{\not{m}}}{\Gamma_0} = -4\bar{f}_\sigma N^{(1)}_{\not{m}} - 4(1-\bar{f}_\sigma)N^{(1)}_{\not{m}}. \tag{63}$$

Solving the above equations, we find

$$N^{(1)}_m = -p_{\sigma,n}\frac{\delta\Gamma^{(1)}_m}{\Gamma_0} \tag{64}$$

$$N^{(1)}_{\not{m}} = -\frac{1}{2}\frac{\delta\Gamma^{(1)}_{\not{m}}}{\Gamma_0} \tag{65}$$

where the coefficient is defined by

$$p_{\sigma,n} \equiv \frac{2(\bar{\rho}_{\sigma,0}+\bar{\rho}_{r,0})}{(4+\epsilon_n)\bar{\rho}_{\sigma,0}+4\bar{\rho}_{r,0}}. \tag{66}$$

Therefore, the curvature perturbation created by the modulation is given by

$$\begin{aligned}\delta N^{(1)} &\equiv N^{(1)}_m - N^{(1)}_{\not{m}}\\ &= \left(-p_{\sigma,n}+\frac{1}{2}\right)\frac{\delta\Gamma^{(1)}}{\Gamma_0}.\end{aligned} \tag{67}$$

Obviously, generation of the curvature perturbation is possible when $\epsilon_n \neq 0$ (i.e, when two components (ρ_σ and ρ_r) are distinguishable in their scaling relations).

B. f_{NL}

Again, we find for the second order perturbations

$$\begin{aligned}N^{(2)}_m &= p_{\sigma,n}\left(\frac{(4+\epsilon_n)^2\bar{f}_\sigma}{2}+8(1-\bar{f}_\sigma)\right)\left(N^{(1)}_m\right)^2\\ &\quad -p_{\sigma,n}\frac{\left(\delta\Gamma^{(1)}\right)^2+\Gamma_0\delta\Gamma^{(2)}}{\Gamma_0^2}\\ N^{(2)}_{\not{m}} &= \frac{1}{2}\left[8\left(N^{(1)}_{\not{m}}\right)^2-\frac{\left(\delta\Gamma^{(1)}\right)^2+\Gamma_0\delta\Gamma^{(2)}}{\Gamma_0^2}\right].\end{aligned} \tag{68}$$

Using the relations

$$\begin{aligned}N^{(1)}_m &= -p_{\sigma,n}\frac{\delta\Gamma^{(1)}}{\Gamma_0}\\ &= -\frac{2p_{\sigma,n}}{1-2p_{\sigma,n}}\delta N^{(1)}\end{aligned} \tag{69}$$

$$\begin{aligned}N^{(1)}_{\not{m}} &= -\frac{1}{2}\frac{\delta\Gamma^{(1)}}{\Gamma_0}\\ &= -\frac{1}{1-2p_{\sigma,n}}\delta N^{(1)},\end{aligned} \tag{70}$$

and the definition (47), we find

$$\begin{aligned}N^{(2)}_m &= \left[\left\{2(4+\epsilon_n)^2\bar{f}_\sigma+32(1-\bar{f}_\sigma)\right\}\frac{p^3_{\sigma,n}}{(1-2p_{\sigma,n})^2}\right.\\ &\quad \left.-\frac{4p_{\sigma,n}(1+A)}{(1-2p_{\sigma,n})^2}\right]\left(\delta N^{(1)}\right)^2\\ N^{(2)}_{\not{m}} &= \left[\frac{4}{(1-2p_{\sigma,n})^2}-\frac{2(1+A)}{(1-2p_{\sigma,n})^2}\right]\left(\delta N^{(1)}\right)^2.\end{aligned} \tag{71}$$

We thus find

$$\begin{aligned}f_{NL} &= \frac{5}{3}\left[\frac{p^3_\sigma(4+\epsilon_n)^2\bar{f}_\sigma+16p^3_\sigma(1-\bar{f}_\sigma)-2p_\sigma-1}{(1-2p_\sigma)^2}\right]\\ &\quad +\frac{5A}{3}\frac{1}{1-2p_\sigma},\end{aligned} \tag{72}$$

where the last term gives the model-dependent contribution. The single-component limit is given by $\bar{f}_\sigma \to 1$ and $p_\sigma \to 2/(4+\epsilon_n)$, where we find significant non-Gaussianity

$$f_{NL} = -\frac{5(4+\epsilon_n)}{3\epsilon_n}-5A, \tag{73}$$

which shows that *the sign of the first term (model-independent contribution) is determined by* ϵ_n*. We find positive sign for* $\epsilon_n < 0$*, while it goes negative when* $\epsilon_n > 0$*. Moreover, the result shows significant non-Gaussianity in the model-independent part, which could be very large even if the decaying component is dominating the Universe.*

In the opposite limit, $\bar{f}_\sigma \to 0$ and $p_\sigma \to 1/2$, we find

$$f_{NL} \propto \frac{1}{(1-2p_\sigma)^2} \gg 1, \tag{74}$$

as expected.

IV. PARTIAL DECAY

More practically, there could be a moment when a fraction of the matter component decays modulated and the decaying component does not have significant interacttion with the other (matter) component. This could be realized when the non-relativistic matter contains particles that belong to the hidden sector.

For the multi-component Universe that contains both matter and radiation, the uniform density hypersurfaces defined for the partial decay is given by

$$\rho_{\sigma,\Gamma} + \Delta\rho_{\sigma,\Gamma} + \rho_{r,\Gamma} \;\equiv\; 3M_p^2\Gamma^2, \tag{75}$$

where the decaying component is denoted by $\Delta\rho_{\sigma,\Gamma}$.

Ignoring isocurvature perturbations and $\delta N_{\rm ini}$, we find

$$\begin{aligned} \bar{f}_\sigma e^{-3N_m} + \bar{f}_\Delta e^{-3N_m} + (1 - f_\sigma - \bar{f}_\Delta)e^{-4N_m} &= \frac{\Gamma^2}{\Gamma_0^2} \\ \bar{f}_\sigma e^{-3N_{\not m}} + \bar{f}_\Delta e^{-4N_{\not m}} + (1 - f_\sigma - \bar{f}_\Delta)e^{-4N_m} &= \frac{\Gamma^2}{\Gamma_0^2}, \end{aligned} \tag{76}$$

where the coefficients are defined by

$$\bar{f}_\sigma \;\equiv\; \frac{\bar{\rho}_{\sigma,0}}{\bar{\rho}_{\sigma,0} + \Delta\bar{\rho}_{\sigma,0} + \bar{\rho}_{r,0}} \tag{77}$$

$$\bar{f}_\Delta \;\equiv\; \frac{\Delta\bar{\rho}_{\sigma,0}}{\bar{\rho}_{\sigma,0} + \Delta\bar{\rho}_{\sigma,0} + \bar{\rho}_{r,0}}. \tag{78}$$

A. First order

We find at first order

$$\begin{aligned} 2\frac{\delta\Gamma^{(1)}}{\Gamma_0} &= -3\bar{f}_\sigma N_m^{(1)} - 3\bar{f}_\Delta N_m^{(1)} \\ &\quad -4(1 - \bar{f}_\sigma - \bar{f}_\Delta)N_m^{(1)} \end{aligned} \tag{79}$$

$$\begin{aligned} 2\frac{\delta\Gamma^{(1)}}{\Gamma_0} &= -3\bar{f}_\sigma N_m^{(1)} - 4\bar{f}_\Delta N_m^{(1)} \\ &\quad -4(1 - \bar{f}_\sigma - \bar{f}_\Delta)N_m^{(1)} \end{aligned} \tag{80}$$

Solving the above equations, we find

$$N_m^{(1)} \;=\; -p_\Delta \frac{\delta\Gamma^{(1)}}{\Gamma_0} \tag{81}$$

$$N_{\not m}^{(1)} \;=\; -\not p_\Delta \frac{\delta\Gamma^{(1)}}{\Gamma_0}, \tag{82}$$

where the coefficients are defined by

$$p_\Delta \;\equiv\; \frac{2(\bar{\rho}_{\sigma,0} + \Delta\bar{\rho}_{\sigma,0} + \bar{\rho}_{r,0})}{3\bar{\rho}_{\sigma,0} + 3\Delta\bar{\rho}_{\sigma,0} + 4\bar{\rho}_{r,0}} \tag{83}$$

$$\not p_\Delta \;\equiv\; \frac{2(\bar{\rho}_{\sigma,0} + \Delta\bar{\rho}_{\sigma,0} + \bar{\rho}_{r,0})}{3\bar{\rho}_{\sigma,0} + 4\Delta\bar{\rho}_{\sigma,0} + 4\bar{\rho}_{r,0}}. \tag{84}$$

Therefore, the curvature perturbation created by the modulation is

$$\begin{aligned} \delta N^{(1)} &\equiv N_m^{(1)} - N_{\not m}^{(1)} \\ &= (-p_\Delta + \not p_\Delta)\frac{\delta\Gamma^{(1)}}{\Gamma_0} \\ &\simeq -p_\Delta r_\Delta \frac{\delta\Gamma^{(1)}}{\Gamma_0}, \end{aligned} \tag{85}$$

where the last approximation is valid when $r_\Delta \ll 1$. Here the coefficient is defined by

$$r_\Delta \equiv \frac{\Delta\bar{\rho}_{\sigma,0}}{3\bar{\rho}_{\sigma,0} + 3\Delta\bar{\rho}_{\sigma,0} + 4\bar{\rho}_{r,0}}. \tag{86}$$

B. Second order

We find for the second order perturbations

$$\begin{aligned} N_m^{(2)} &= p_\Delta\left(8 - \frac{7}{2}(\bar{f}_\sigma + \bar{f}_\Delta)\right)\left(N_m^{(1)}\right)^2 \\ &\quad -p_\Delta \frac{\left(\delta\Gamma^{(1)}\right)^2 + \Gamma_0\delta\Gamma^{(2)}}{\Gamma_0^2} \end{aligned} \tag{87}$$

$$\begin{aligned} N_{\not m}^{(2)} &= \not p_\Delta\left(8 - \frac{7}{2}\bar{f}_\sigma\right)\left(N_m^{(1)}\right)^2 \\ &\quad -\not p_\Delta \frac{\left(\delta\Gamma^{(1)}\right)^2 + \Gamma_0\delta\Gamma^{(2)}}{\Gamma_0^2}. \end{aligned} \tag{88}$$

Using the relations

$$\begin{aligned} N_m^{(1)} &= -p_\Delta \frac{\delta\Gamma^{(1)}}{\Gamma_0} \\ &= \frac{p_\Delta}{p_\Delta - \not p_\Delta}\delta N^{(1)} \end{aligned} \tag{89}$$

$$\begin{aligned} N_{\not m}^{(1)} &= -\not p_\Delta \frac{\delta\Gamma^{(1)}}{\Gamma_0} \\ &= \frac{\not p_\Delta}{p_\Delta - \not p_\Delta}\delta N^{(1)}, \end{aligned} \tag{90}$$

and the definition (47), we find

$$\begin{aligned} N_m^{(2)} &= \left[\frac{p_\Delta^3(16 - 7\bar{f}_\sigma - 7\bar{f}_\Delta)}{2(p_\Delta - \not p_\Delta)^2} - \frac{p_\Delta(1+A)}{(p_\Delta - \not p_\Delta)^2}\right]\left(\delta N^{(1)}\right)^2 \\ N_{\not m}^{(2)} &= \left[\frac{\not p_\Delta^3(16 - 7\bar{f}_\sigma)}{2(p_\Delta - \not p_\Delta)^2} - \frac{\not p_\Delta(1+A)}{(p_\Delta - \not p_\Delta)^2}\right]\left(\delta N^{(1)}\right)^2. \end{aligned} \tag{91}$$

Then, f_{NL} is calculated from

$$f_{NL} \;=\; \frac{5}{6}\frac{N_m^{(2)} - N_{\not m}^{(2)}}{\left(\delta N^{(1)}\right)^2}. \tag{92}$$

V. CONCLUSION AND DISCUSSION

The early Universe after inflation may have many components. They could be oscillations, topological defects, relativistic and non-relativistic particles. If those components are decaying into radiation, there could be generation of the curvature perturbations. In this paper, the mechanism of the modulated decay has been considered for the multi-component Universe.

In this paper we found the basic formulation, which is useful in calculating modulated decays in the multi-component Universe.

We found useful results, in which the non-Gaussianity parameter is separated into model-dependent and independent terms. Interestingly, the sign of the model-independent f_{NL} may flip depending on the scaling of the decaying component.

We find that the component whose scaling is very close to the radiation ($\epsilon_n \simeq 0$) could generate significant non-Gaussianity in the single-component limit. Importantly, the sign could be either positive or negative. In that way, the modulated decay of the oscillations may *crucially depend on the amplitude* at the decay. For instance, consider the potential for the oscillations given by

$$V(\sigma) \simeq \frac{1}{2}m^2\sigma^2 + \frac{\lambda_4}{4}\sigma^4 + \frac{\lambda_6}{6}\frac{\varphi^6}{M_p^2}. \tag{93}$$

If the oscillations decay when φ^6 is dominant, one will find $f_{NL} < 0$. If the oscillations decay when φ^4 is dominant, one will find $|f_{NL}| \gg 1$, where the sign could be either positive or negative. The scaling of the density changes during the oscillations. One will find conventional result when the quadratic term is dominating. In the intermediate region one may find the density scaling $\rho_\sigma \propto a^{-k}$, where (effectively) $3 \le k \le 6$ is possible. As the result, in the practical calculations the curvature perturbations and the non-Gaussianity may depend crucially on the amplitude of the oscillations.

Note added: While finalizing this paper, we found a couple of papers [24] which has some overlaps with our models.

VI. ACKNOWLEDGMENT

T.M wishes to thank K. Shima for encouragement, and his colleagues at Nagoya university and Lancaster university for their kind hospitality and many invaluable discussions. This work is supported in part by Grant-in-Aid for Scientific research from the Ministry of Education, Sci- ence, Sports, and Culture (MEXT), Japan, No. 21111006, No. 22244030, and No. 23540327 (K.K.). S.E. is supported by the Grant-in-Aid for Nagoya University Global COE Program, "Quest for Fundamental Principles in the Universe: from Particles to the Solar System and the Cosmos".

Appendix A: Non-Linear formalism and the curvaton mechanism

In this section we first review the basics of the non-linear formalism and then compare our results with Ref.[24]. Notations in Ref.[24] are discriminated by the subscripts "LT-A".

The non-linear formalism in the curvaton mechanism is given by the formula

$$\zeta_\sigma \;=\; \delta N + \frac{1}{3}\ln\left(\frac{\rho_\sigma}{\bar\rho_\sigma}\right), \tag{A1}$$

$$\zeta_r \;=\; \delta N + \frac{1}{4}\ln\left(\frac{\rho_r}{\bar\rho_r}\right). \tag{A2}$$

Here δN is the perturbation of N, which is measured between two hypersurfaces. Usually the one is the flat hypersurface and the other is a uniform density hypersurface. Besides δN, we have to define the other quantities (ρ_σ, ρ_r) and ($\bar\rho_\sigma$, $\bar\rho_r$). Following the definition of δN, those quantities are defined on the uniform density hypersurface on which δN is defined.

We thus find at the uniform density hypersurface $H = H_A$;

$$\zeta_{\sigma,A} \;=\; \delta N_A(t_A) + \frac{1}{3}\ln\left(\frac{\rho_{\sigma,A}(x,t_A)}{\bar\rho_{\sigma,A}(t_A)}\right), \tag{A3}$$

$$\zeta_{r,A} \;=\; \delta N_A(t_A) + \frac{1}{4}\ln\left(\frac{\rho_{r,A}(x,t_A)}{\bar\rho_{r,A}(t_A)}\right). \tag{A4}$$

Solving these equations we find

$$\begin{aligned} \rho_{\sigma,A} &= \bar\rho_{\sigma,A}e^{3(\zeta_{\sigma,A}-\delta N_A)}, \\ \rho_{r,A} &= \bar\rho_{r,A}e^{4(\zeta_{r,A}-\delta N_A)}. \end{aligned} \tag{A5}$$

The trivial identity is

$$\frac{\rho_{\sigma,A}+\rho_{r,A}}{\bar\rho_{\sigma,A}+\bar\rho_{r,A}} = 1, \tag{A6}$$

where $\rho_{\sigma,A}$ and $\rho_{r,A}$ can be replaced using Eq.(A5). We find the equation

$$\bar f_{\sigma,A}e^{3(\zeta_{\sigma,A}-\delta N_A)} + (1-\bar f_{\sigma,A})e^{4(\zeta_{r,A}-\delta N_A)} \;=\; 1, \tag{A7}$$

where the ratio is defined by

$$\bar f_{\sigma,A} \;=\; \frac{\bar\rho_{\sigma,A}}{\bar\rho_{\sigma,A}+\bar\rho_{r,A}}. \tag{A8}$$

We find at first order

$$\begin{aligned} \delta N_A &= r_{\sigma,A}\zeta_{\sigma,A} + (1-r_{\sigma,o})\zeta_{r,o} \\ &= \delta N_A \\ &\quad + \frac{r_A}{3}\ln\left(\frac{\rho_{\sigma,A}}{\bar\rho_{\sigma,A}}\right) + \frac{(1-r_A)}{4}\ln\left(\frac{\rho_{r,A}}{\bar\rho_{r,A}}\right). \end{aligned} \tag{A9}$$

The trivial identity is

$$\frac{r_A}{3}\ln\left(\frac{\rho_{\sigma,A}}{\bar\rho_{\sigma,A}}\right) + \frac{(1-r_A)}{4}\ln\left(\frac{\rho_{r,A}}{\bar\rho_{r,A}}\right) \;=\; 0. \tag{A10}$$

For the expansion $\delta\rho_i \equiv \rho_i - \bar\rho_i$, the above equation gives the obvious identity

$$\delta\rho_{\sigma,A} + \delta\rho_{r,A} \;=\; 0. \tag{A11}$$

One may evaluate the non-linear formalism **away** from $H = H_A$. (See Fig.3.) Choosing **another** hypersurface at $H = H_B$, one can evaluate

$$\begin{aligned} \delta N_B &= r_{\sigma,B}\zeta_{\sigma,B} + (1-r_{\sigma,B})\zeta_{r,B} \\ &= r_{\sigma,B}\zeta_{\sigma,A} + (1-r_{\sigma,B})\zeta_{r,A}, \end{aligned} \tag{A12}$$


δN_A
$H = H_A$
$(\rho_{\sigma,A}, \rho_{r,A}); (\bar{\rho}_{\sigma,A}, \bar{\rho}_{r,A})$
$\delta N_{\rm curv}$
δN_B
$H = H_B$
$(\rho_{\sigma,B}, \rho_{r,B}); (\bar{\rho}_{\sigma,B}, \bar{\rho}_{r,B})$

FIG. 3: Component perturbations are defined on the uniform density hypersurfaces. For instance, $\zeta_{\sigma,A} \equiv \delta N_A + \frac{1}{3}\ln\left(\frac{\rho_{\sigma,A}}{\bar{\rho}_{\sigma,A}}\right)$ is defined using δN_A (measured from the flat hypersurfaces to $H = H_A$), $\rho_{\sigma,A}$ and its mean value $\bar{\rho}_{\sigma,A}$ (both are defined on $H = H_A$). Note that the isocurvature inhomogeneity $\delta\rho_{\sigma,A}(x,t_A) \equiv \rho_{\sigma,A}(x,t_A) - \bar{\rho}_{\sigma,A}(t_A)$ is possible in the uniform density hypersurfaces when the Universe is multi-component.

where the constancy of the component perturbations $\zeta_{i,A} = \zeta_{i,B}$ has been used.

Note that $\delta N_{\rm curv} \equiv \delta N_B - \delta N_A$ gives the "evolution" between the two hypersurfaces. We thus find for $r_{\sigma,B} \gg r_{\sigma,A}$:

$$\begin{aligned}\delta N_{\rm curv} &= (r_{\sigma,B} - r_{\sigma,A})\zeta_{\sigma,A} - (r_{\sigma,B} - r_{\sigma,A})\zeta_{r,A} \\ &\simeq r_{\sigma,B}\left[\frac{\delta\rho_{\sigma,A}}{3\bar{\rho}_{\sigma,A}}\right]. \end{aligned} \tag{A13}$$

Note that δN_A does not appear in the final result because of the obvious cancellation.

If one uses H_A at the beginning of the curvaton oscillation and H_B at the decay, $\delta N_{\rm curv}$ gives the evolution of the curvature perturbation in the conventional curvaton mechanism.

The curvature perturbation generated before the curvaton oscillation can be included as $\delta N_{\rm inf} \simeq \delta N_A$. Then one will find the post-decay curvature perturbation

$$\delta N_B \simeq \delta N_{\rm inf} + \delta N_{\rm curv}. \tag{A14}$$

1. Why difficult?

The formalism that can be applied for the modulation at the end has already been discussed by Enomoto-Kohri-Matsuda(EKM) in Ref.[11]. When the decay is modulated, the non-linear formalism at the end can be given by

$$\begin{aligned}\zeta_{\sigma,\rm E} &= \delta N_{\rm E} + \frac{1}{3}\ln\left(\frac{\rho_{\sigma,d}}{\bar{\rho}_\sigma}\right) + \frac{1}{4}\ln\left(\frac{\rho_{\sigma,\rm E}}{\rho_{\sigma,d}}\right) \\ &= \delta N_{\rm E} + \frac{1}{3}\ln\left(\frac{\rho_{\sigma,\rm E}}{\bar{\rho}_\sigma}\right) + \Delta_E \\ \zeta_{r,\rm E} &= \delta N_{\rm E} + \frac{1}{4}\ln\left(\frac{\rho_{r,\rm E}}{\bar{\rho}_r}\right), \end{aligned} \tag{A15}$$

where we defined

$$\Delta_{\rm E} \equiv \frac{1}{12}\ln\left(\frac{\rho_{\sigma,d}}{\rho_{\sigma,\rm E}}\right). \tag{A16}$$

Here the subscript "E" means that the quantities are evaluated at $t = t_{\rm E}$, which denotes the time after the decay. The quantities denoted by "bar" is placed before the decay.

Again, the trivial identity

$$\frac{\rho_{\sigma,\rm E} + \rho_{r,\rm E}}{\bar{\rho}_\sigma + \bar{\rho}_r} = \frac{H_{\rm E}^2}{\bar{H}^2} \tag{A17}$$

gives

$$\bar{f}_\sigma e^{3(\zeta_{\sigma,\rm E} - \delta N_{\rm E} - \Delta_{\rm E})} + (1 - \bar{f}_\sigma)e^{4(\zeta_r - \delta N_{\rm E})} = \frac{H_{\rm E}^2}{\bar{H}^2}, \tag{A18}$$

It is possible to identify $H_E \equiv \Gamma$ and $\bar{H} \equiv \Gamma_0$ to find $\Delta_E \equiv 0$, however in that case the perturbation of $H_E = \Gamma$ is a problem. We thus need some trick to find the exact result.

At this moment we have two solutions for the problem. One is discussed in this paper, and the other is discussed in Ref.[24] by two groups. For instance, Langlois and Takahashi introduced a new parameter δN_D to define

$$\zeta_\sigma = \delta N_D + \frac{1}{3}\ln\left(\frac{\rho_{\sigma,d}}{\bar{\rho}_\sigma}\right), \tag{A19}$$

where δN_D *cannot be identified with the curvature perturbation* ζ*, while* ζ_σ *is identical to the conventional component pertubation.* Below, we will take a closer look at these definitions.

2. How to understand the calculation of Ref.[24] in terms of our set-ups

We are going to show the obvious correspondences between the quantities defined in Ref.[24] and our set-ups presented in Fig.2.

In Ref.[24], they have defined the non-linear formalism

$$\zeta_\sigma = \delta N_D + \frac{1}{3}\ln\left(\frac{\rho_\sigma(t_D)}{\bar{\rho}_\sigma(\bar{t}_D)}\right), \tag{A20}$$

where the uniform density hypersurfaces defined by t_D ($H = \Gamma$) and $\bar{t}_D$ ($H = \bar{\Gamma}$) are not identical. δN_D is defined for the hypersurface $H = \Gamma$, but it is not the usual curvature perturbation. (See Fig.4.)



FIG. 4: Unlike Fig.3, component perturbations of ref.[24] are defined using multiple hypersurfaces. For instance, $\zeta_\sigma|_{(LT-A)} \equiv \delta N_D + \frac{1}{3}\ln\left(\frac{\rho_\sigma(t_D)}{\bar{\rho}_{\sigma,A}(t_D)}\right)$ is defined using both $H=\bar{\Gamma}$ and $H=\Gamma$. As a result, $-\frac{1}{2}\ln(1+\delta_\Gamma)$ has to be subtracted from δN_D to get the curvature perturbation ζ.

In our formalism, these quantities are corresponding to

$$\begin{aligned}
\delta N_D|_{(LT-A)} &\leftrightarrow \delta N_{\rm ini} + N_m \\
\rho_\sigma(t_D)|_{(LT-A)} &\leftrightarrow \rho_{\sigma,\Gamma,m} = \rho_{\sigma,0}e^{-3N_m} \\
\bar{\rho}_\sigma(\bar{t}_D)|_{(LT-A)} &\leftrightarrow \bar{\rho}_{\sigma,0}.
\end{aligned} \tag{A21}$$

We thus find

$$\zeta_\sigma|_{(LT-A)} \leftrightarrow \zeta_\sigma. \tag{A22}$$

They also defined

$$(1+\delta_\Gamma)^2 \equiv \Gamma^2/\bar{\Gamma}^2, \tag{A23}$$

which gives the correspondence

$$(1+\delta_\Gamma)^2|_{(LT-A)} \leftrightarrow e^{-4N_{\not m}}. \tag{A24}$$

Finally, they have defined the post-decay curvature perturbation

$$\zeta = \delta N_D + \frac{1}{2}\ln(1+\delta_\Gamma), \tag{A25}$$

which corresponds to

$$\zeta|_{(LT-A)} \leftrightarrow \delta N_{\rm ini} + N_m - N_{\not m}. \tag{A26}$$

In our calculation the curvature perturbation generated by the modulated decay is given by

$$\delta N_{\rm mod} \equiv N_m - N_{\not m}. \tag{A27}$$

Therefore, the correspondence is obvious between our calculation and Ref.[24].

However, in finding the curvaton contribution they evaluated

$$\zeta = \zeta_r - \frac{r}{6}\delta_\Gamma + \frac{r}{3}S, \tag{A28}$$

where the first and the last terms are originally given by

$$\zeta_r + \frac{r}{3}S = r\zeta_\sigma + (1-r)\zeta_r \equiv \delta N_{\rm ini}, \tag{A29}$$

where $\delta N_{\rm ini}$ is defined in this paper. If the curvaton hypothesis is valid and the component perturbations are constant, one may evaluate the component perturbations at $H = H_{\rm osc}$:

$$\zeta_\sigma = \zeta_\sigma(t_{\rm osc}) = \delta N_{\rm inf} + \frac{1}{3}\ln\frac{\rho_\sigma(t_{\rm osc})}{\bar{\rho}_\sigma(t_{\rm osc})}, \tag{A30}$$

where $\delta N_{\rm inf}$ denotes the curvature perturbation just at the beginning of the oscillation. Substituting the component perturbations (defined at $H_{\rm osc}$) into the above equation, one finds

$$\begin{aligned}
\zeta_r + \frac{r}{3}S &= r\zeta_\sigma(t_{\rm osc}) + (1-r)\zeta_r(t_{\rm osc}) \\
&= \delta N_{\rm inf} + \frac{r}{3}\ln\frac{\rho_\sigma(t_{\rm osc})}{\bar{\rho}_\sigma(t_{\rm osc})} \\
&\quad + \frac{1-r}{4}\ln\frac{\rho_r(t_{\rm osc})}{\bar{\rho}_r(t_{\rm osc})}.
\end{aligned} \tag{A31}$$

Although trivial deformation is needed, *it is easy to find that the result is consistent with Eq.(A13) without using approximations.*

Using the above formula, they started perturbation with regard to the perturbation of S. For instance, Langlois and Takahashi considered for the "curvaton perturbation"

$$S \equiv 3(\zeta_\sigma - \zeta_r) = \frac{2\sigma}{\sigma} - \frac{\delta\sigma^2}{\sigma^2} + \frac{2}{3}\frac{\delta\sigma^3}{\sigma^3}, \tag{A32}$$

and for the "inflaton perturbation"

$$\zeta_r = \frac{H}{\dot{\phi}}\delta\phi \simeq \delta N_{\rm inf}. \tag{A33}$$

These definitions (in fact they are approximations valid only when one can disregard $\delta\rho_r/\rho_r$) are based on the usual curvaton hypothesis.

Although in this paper we were not focusing on the mixed perturbations, our formalism is carefully prepared to explain the modulated decay in the multi-component Universe, where the curvaton hypothesis could not apply.

---

[1] D. H. Lyth, A. R. Liddle, "The primordial density perturbation: Cosmology, inflation and the origin of structure," Cambridge, UK: Cambridge Univ. Pr. (2009) 497 p.

[2] D. Wands, K. A. Malik, D. H. Lyth and A. R. Liddle, "A New approach to the evolution of cosmological perturbations on large scales," Phys. Rev. D **62**, 043527 (2000) [astro-ph/0003278]; T. Matsuda, "Delta-N formalism for the evolution of the curvature perturbations in generalized multi-field inflation," Phys. Lett. B **682**, 163 (2009) [arXiv:0906.2525 [hep-th]]; T. Matsuda, "Elliptic Inflation: Generating the curvature perturbation without slow-roll," JCAP **0609**, 003 (2006) [hep-ph/0606137].

[3] F. Bernardeau and J. -P. Uzan, "NonGaussianity in multifield inflation," Phys. Rev. D **66**, 103506 (2002) [hep-ph/0207295]; L. E. Allen, S. Gupta and D. Wands, "Non-gaussian perturbations from multi-field inflation," JCAP **0601**, 006 (2006) [astro-ph/0509719]; H. Assadullahi, H. Firouzjahi, M. H. Namjoo and D. Wands, "Curvaton and the inhomogeneous end of inflation," arXiv:1207.7006 [astro-ph.CO].

[4] T. Matsuda, "Modulated Inflation," Phys. Lett. B **665**, 338 (2008) [arXiv:0801.2648 [hep-ph]]; T. Matsuda, "Modulated inflation from kinetic term," JCAP **0805**, 022 (2008) [arXiv:0804.3268 [hep-th]].

[5] T. Moroi, T. Takahashi, "Effects of cosmological moduli fields on cosmic microwave background," Phys. Lett. **B522**, 215-221 (2001). [hep-ph/0110096]. D. H. Lyth, D. Wands, "Generating the curvature perturbation without an inflaton," Phys. Lett. **B524**, 5-14 (2002). [hep-ph/0110002].

[6] K. Dimopoulos, K. Kohri and T. Matsuda, "The hybrid curvaton," Phys. Rev. D **85**, 123541 (2012) [arXiv:1201.6037 [hep-ph]]; T. Matsuda, "Free light fields can change the predictions of hybrid inflation," JCAP **1204**, 020 (2012) [arXiv:1204.0303 [hep-ph]].

[7] K. Dimopoulos, K. Kohri, D. H. Lyth and T. Matsuda, "The inflating curvaton," JCAP **1203**, 022 (2012) [arXiv:1110.2951 [astro-ph.CO]];

[8] K. Kohri, C. -M. Lin and T. Matsuda, "PBH from the inflating curvaton," arXiv:1211.2371 [hep-ph].

[9] G. Dvali, A. Gruzinov and M. Zaldarriaga, "A new mechanism for generating density perturbations from inflation," Phys. Rev. D **69**, 023505 (2004) [arXiv:astro-ph/0303591].

[10] D. H. Lyth, "Generating the curvature perturbation at the end of inflation," JCAP **0511**, 006 (2005) [astro-ph/0510443]; F. Bernardeau, L. Kofman and J. -P. Uzan, "Modulated fluctuations from hybrid inflation," Phys. Rev. D **70**, 083004 (2004) [astro-ph/0403315].

[11] S. Enomoto, K. Kohri and T. Matsuda, "Non-Gaussianity in the unified curvaton mechanism : The generalized curvaton mechanism that comprehends modulation at the transition," arXiv:1210.7118 [hep-ph].

[12] T. Matsuda, "Cosmological perturbations from inhomogeneous preheating and multi-field trapping," JHEP **0707**, 035 (2007) [arXiv:0707.0543 [hep-th]].

[13] T. Matsuda, "Cosmological perturbations from an inhomogeneous phase transition," Class. Quant. Grav. **26**, 145011 (2009). [arXiv:0902.4283 [hep-ph]]; M. Kawasaki, T. Takahashi and S. Yokoyama, "Density Fluctuations in Thermal Inflation and Non-Gaussianity," JCAP **0912**, 012 (2009) [arXiv:0910.3053 [hep-th]].

[14] G. N. Felder, L. Kofman and A. D. Linde, Phys. Rev. D **60**, 103505 (1999) [hep-ph/9903350]; T. Matsuda, JCAP **0708**, 003 (2007) [arXiv:0707.1948 [hep-ph]].

[15] T. Matsuda, "Topological curvatons," Phys. Rev. D **72**, 123508 (2005) [hep-ph/0509063].

[16] D. H. Lyth, "Can the curvaton paradigm accommodate a low inflation scale?," Phys. Lett. B **579** (2004) 239 [hep-th/0308110].

[17] K. A. Malik and D. H. Lyth, "A numerical study of non-gaussianity in the curvaton scenario," JCAP **0609**, 008 (2006) [astro-ph/0604387].

[18] M. Zaldarriaga, "Non-Gaussianities in models with a varying inflaton decay rate," Phys. Rev. D **69**, 043508 (2004) [astro-ph/0306006]; T. Suyama and M. Yamaguchi, "Non-Gaussianity in the modulated reheating scenario," Phys. Rev. D **77** (2008) 023505 [arXiv:0709.2545 [astro-ph]]; K. Kohri, D. H. Lyth and C. A. Valenzuela-Toledo, "Preheating and the non-gaussianity of the curvature perturbation," JCAP **1002**, 023 (2010) [Erratum-ibid. **1009**, E01 (2011)] [arXiv:0904.0793 [hep-ph]]; M. Cicoli, G. Tasinato, I. Zavala, C. P. Burgess and F. Quevedo, "Modulated Reheating and Large Non-Gaussianity in String Cosmology," JCAP **1205**, 039 (2012) [arXiv:1202.4580 [hep-th]].

[19] K. -Y. Choi and O. Seto, "Modulated reheating by curvaton," arXiv:1204.1419 [astro-ph.CO].

[20] D. H. Lyth, K. A. Malik and M. Sasaki, "A General proof of the conservation of the curvature perturbation," JCAP **0505**, 004 (2005) [astro-ph/0411220];

[21] D. Langlois, F. Vernizzi and D. Wands, "Non-linear isocurvature perturbations and non-Gaussianities," JCAP **0812**, 004 (2008) [arXiv:0809.4646 [astro-ph]].

[22] M. Sasaki, J. Valiviita and D. Wands, "Non-Gaussianity of the primordial perturbation in the curvaton model," Phys. Rev. D **74**, 103003 (2006) [astro-ph/0607627].

[23] M. S. Turner, "Coherent Scalar Field Oscillations in an Expanding Universe," Phys. Rev. D **28**, 1243 (1983).

[24] D. Langlois and T. Takahashi, "Density Perturbations from Modulated Decay of the Curvaton," arXiv:1301.3319 [astro-ph.CO]; H. Assadullahi, H. Firouzjahi, M. H. Namjoo and D. Wands, "Modulated curvaton decay," arXiv:1301.3439 [hep-th].